An Adaptive Optics Survey for Companions to Stars with Extra-Solar Planets

J. P. Lloyd, M. C. Liu, J. R. Graham, M. Enoch, P. Kalas, G. W. Marcy, D. Fischer, J. Patience, B. Macintosh, D. T. Gavel, S. S. Olivier, C. E. Max, R. White, A. M. Ghez, and I. S. McLean

This article was submitted to Planetary Systems in the Universe: Observation, Formation and Evolution International Astronomical Union Symposium 2000, Manchester, England, August 7-11, 2000

U.S. Department of Energy



November 27, 2000

DISCLAIMER

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

This is a preprint of a paper intended for publication in a journal or proceedings. Since changes may be made before publication, this preprint is made available with the understanding that it will not be cited or reproduced without the permission of the author.

This report has been reproduced directly from the best available copy.

Available electronically at http://www.doc.gov/bridge

Available for a processing fee to U.S. Department of Energy
And its contractors in paper from
U.S. Department of Energy
Office of Scientific and Technical Information
P.O. Box 62
Oak Ridge, TN 37831-0062

Telephone: (865) 576-8401 Facsimile: (865) 576-5728 E-mail: reports@adonis.osti.gov

Available for the sale to the public from U.S. Department of Commerce National Technical Information Service 5285 Port Royal Road Springfield, VA 22161 Telephone: (800) 553-6847

Facsimile: (703) 605-6900 E-mail: orders@ntis.fedworld.gov

Online ordering: http://www.ntis.gov/ordering.htm

OR

Lawrence Livermore National Laboratory
Technical Information Department's Digital Library
http://www.llnl.gov/tid/Library.html

An Adaptive Optics Survey for Companions to Stars with Extra-Solar Planets

James P. Lloyd, Michael C. Liu, James R. Graham, Melissa Enoch, Paul Kalas, Geoffrey W. Marcy, Debra Fischer

Department of Astronomy, University of California, Berkeley, CA 94720. USA

Jennifer Patience, Bruce Macintosh, Donald T. Gavel, Scot S. Olivier, Claire E. Max

Institute of Geophysics and Planetary Physics, Lawrence Livermore National Laboratory, Livermore, CA 95064, USA

Russel White, Andrea M. Ghez, Ian S. McLean

Department of Physics and Astronomy, University of California, Los Angeles, CA 90095, USA

Abstract. We have undertaken an adaptive optics imaging survey of extrasolar planetary systems and stars showing interesting radial velocity trends from high precision radial velocity searches. Adaptive Optics increases the resolution and dynamic range of an image, substantially improving the detectability of faint close companions. This survey is sensitive to objects less luminous than the bottom of the main sequence at separations as close as 1". We have detected stellar companions to the planet bearing stars HD 114762 and Tau Boo. We have also detected a companion to the non-planet bearing star 16 Cyg A.

1. Introduction

The Lick Adaptive Optics System is a Shack-Hartmann Laser Guide star AO system on the Lick Observatory Shane 3m telescope at Mt Hamilton, California (Max et al. 1997). For these observations, we used the system in natural guide star mode, with the bright star serving as wavefront reference. Under good conditions, the system produces diffraction limited images (0".15 FWHM) with a strehl ratio of 0.7 in the K band (2.2 μ m).

The AO system feeds a 256×256 pixel infrared camera, IRCAL (Lloyd et al. 2000), which reimages the field of view at 0″.076 per pixel. The camera incorporates a cold focal plane with an occulting finger to obtain high dynamic range images. For this program, we typically take a few minutes of integration of unsaturated images to obtain coverage close to the star, and deep exposures in coronagraphic mode to detect faint companions at larger separations.

We have selected targets from those stars with radial velocity planets, or with radial velocity trends from the Lick and Keck radial velocity surveys.

2. Results

HD 155423 (see Fig 1) shows substantial scatter in precision radial velocity measurements. High resolution imaging shows HD 155423 is a hierarchical triple, with a 0"2 close binary (8AU projected separation) M3 dwarf pair, separated 1"5 from the F8 dwarf primary.

A faint object was previously discovered near 16 Cyg A, but was not known to be physically associated (Hauser & Marcy 1999). We have confirmed by proper motion measurements that the ΔK =5.4 object is a physically associated M5 dwarf (see Fig 1). Radial velocity measurements show a shallow linear trend.

HD 114762 has a radial velocity companion with an 84 day period and Msin i=11 M_{Jup} (Latham et al. 1989). We have detected a $\Delta K=7.3$ companion 3".3 from the primary (see Fig 1). JHK photometry and follow up Keck AO/NIRSPEC spectroscopy (McLean et al. 2000) reveal the companion to be a late M subdwarf. The physical association of this companion is confirmed by proper motion over a 2 year baseline (Patience et al. 1998).

Radial velocity measurements of HR 6623 show a nearly linear trend over 13 years (see Fig 1). This would classify this object as a poorly determined single lined spectroscopic binary. AO imaging resolves the companion, which is an M5 dwarf. Further radial velocity and astrometric observations will allow accurate mass determinations.

Tau Boo hosts an Msin $i=3.9 \,\mathrm{M}_{Jup}, 3.3$ day period planet, and shows radial velocity residuals (see Fig 1). It has an M2V companion that was discovered in 1849 at 10".3. At present it is at 2".82, with 0".01 per year of orbital motion. Although it has been suggested that there may be additional companions in the system (Wiedemann, Deming, & Bjoraker 2000), we do not detect any additional companions, and attribute the velocity residuals to the stellar companion.

Acknowledgments. This research was supported in part by the National Science Foundation under a cooperative agreement with the Center for Adaptive Optics, Agreement No. AST-987678. Work on the Lick adaptive optics system was performed in part under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory, Contract number W-7405-ENG-48

References

Cochran, W., Hatzes, A., Butler, R. P. & Marcy, G. W. 1997, ApJ, 483, 457 Hauser, H. M. & Marcy, G. W. 1999, PASP, 111, 321 Latham, D., Stefanik, R., Mazeh, T., et al. 1989, Nature, 339, 38 Lloyd, J. P., Liu, M. C., Macintosh, B. A., Severson, S. A., Deich, W. T. S. & Graham J. R. 2000, in Proc. SPIE 4008, Optical and IR Telescope Instrumentation and Detectors, ed. M. Iye & A. F. Moorwood, (Washington: SPIE) Max, C. E., Olivier, S. S., Friedman, H. W., et al., Science, 277, 1649 McLean, I. S. Wilcox, M. K., Becklin, E. E. et al. 2000, ApJ, 533, L45 Patience, J., Ghez, A. M., White, R. J. et al. 1998, BAAS, 193, 9708 Wiedemann, G., Deming, D., Gjoraker, G. 2000, astro-ph/0007216




Figure 1. K band images of stars with detected companions; J band spectra of HD 114762B; Radial Velocity data for HR 6623 and Tau Boo (residual)